

Variance Estimation for Estimators of Between-Year Change in Totals from Two Stratified Bernoulli Samples

Kimberly Henry¹, Valerie Testa¹, and Richard Valliant²

¹Statistics of Income, P.O. Box 2608, Washington DC 20013-2608

²University of Michigan, 1218 Lefrak Hall, College Park MD 20742

Abstract: This paper provides the theoretical framework for estimating the variance of the difference in two years' totals estimated under the stratified Bernoulli sample design. We provide a design-unbiased estimator that takes into account two practical problems: a large overlap of units between two years' samples and "stratum jumpers," which are population and sample units that shift across strata from one year to another. Both problems affect estimating the covariance term in the variance of the difference. The estimator is applied to data from the Statistics of Income Division's individual tax return sample. Naïve variance estimates using only the separate years' variances are compared to show the effect of ignoring the estimated covariance.

Key Words: Horvitz-Thompson estimator, poststratification, survey sampling

1. Introduction and Universe of Tax Returns for Two Years

This paper contains the theoretical background necessary to produce variance estimates of year-to-year changes between totals estimated from the Statistics of Income (SOI) Division's Individual Tax Return sample, a stratified Bernoulli sample. The underlying theory here is modified from the theory in Valliant and Casady (1998). Related work for similar sample designs can be found in Berger (2004), Nordberg (2000), and Wood (2008).

We consider two estimators of the finite population total: the Horvitz-Thompson (HT) and post-stratification (PS) estimators. SOI uses the PS estimator to estimate yearly totals, but both are considered here for comparison purposes. Suppose that the strata are ordered by increasing size of the sampling rate, i.e., the sampling rate for stratum 2 is greater than or equal to the rate for stratum 1, and so on. Both estimators are affected by the location of sample units within strata in both years, so we define the following:

- $U_{h_1 0}$ = returns in stratum h_1 that file only at time t_1 (deaths after time t_1 and before time t_2)
- $U_{0 h_2}$ = returns in stratum h_2 that file only at time t_2 (births after time t_1 and before time t_2)
- $U_{h_1 h_2}$ = returns in stratum h_1 at time t_1 and stratum h_2 at time t_2 that file returns at both times, for $h_1 < h_2$ (units that move to strata with a higher sampling rate in year 2), $h_1 = h_2$ (units that stay in the same strata), or $h_1 > h_2$ (units moving to strata with lower sampling rates in year 2).

Using this notation, the two universes can be partitioned into a 2-way grid based on stratum membership at times t_1 and t_2 , shown in Table 1 on the following page. If the set of strata is the same in the two years, then $H_1 = H_2$ and returns on the diagonal of Table 1 remain in the same stratum between the two years, while the "stratum jumpers" (returns that shift to different strata between the years) lie on the off-diagonal. For sample selection purposes, the stratum h_1 and h_2 universes at times t_1 and t_2 are the union of all units (here tax returns) down column h_2 and across row h_1 of Table 1, respectively:

$$U_{h_1 \bullet} = \bigcup_{h_2=0}^{H_2} U_{h_1 h_2} \quad \text{and} \quad U_{\bullet h_2} = \bigcup_{h_1=0}^{H_1} U_{h_1 h_2} .$$

2. Sample Design

The stratified Bernoulli design is used by most of SOI's cross-sectional studies. In each study's frame population, every unit has a unique identifier - the Social Security Number (SSN) of the primary tax filer in the Individual study and the Employer Identification Number for Corporate and Tax Exempt organizations' tax returns. Each return's unique identifier is used to produce a permanent random number (PRN) between 0 and 1, denoted r_i . For a given year, unit i is selected for a sample if

$$r_i < \pi_h , \tag{2.1}$$

where π_h is the pre-assigned sampling rate for stratum h that tax return i belongs to. Stratification for SOI's Form 1040 sample used various criteria, including size of total gross positive/negative income indexed for inflation and an indicator of the return's "usefulness" for tax policy modeling purposes, to create 208 strata (see Testa and Scali 2005 for details). Every week of the IRS processing year, all 1040, 1040A, and 1040EZ returns in the frame population were assigned to a stratum as

the data were transcribed by IRS. Due to this weekly schedule of available frame data, SOI uses stratified Bernoulli sampling to select its samples. Similar procedures are used by other tax agencies (e.g., Revenue Canada, Cooney 2008).

Table 1: Partition of Universe at Two Times

	Time t_2 Stratum Membership				
Time t_1 Stratum Membership	0 (deaths in t_1)	1	...	H_2	Stratum universe at time t_1
0 (births in t_2)	--	U_{01}	...	U_{0H_2}	--
1	U_{10}	U_{11}	...	U_{1H_2}	$U_{1\bullet} = \bigcup_{h_2=0}^{H_2} U_{1h_2}$
\vdots	\vdots	\vdots	\ddots	\vdots	\vdots
H_1	U_{H_10}	U_{H_11}	...	$U_{H_1H_2}$	$U_{H_1\bullet} = \bigcup_{h_2=0}^{H_2} U_{H_1h_2}$
Stratum universe at time t_2	--	$U_{\bullet 1} = \bigcup_{h_1=0}^{H_1} U_{h_11}$...	$U_{\bullet H_2} = \bigcup_{h_1=0}^{H_1} U_{h_1H_2}$	

SOI’s Individual sample consisted of two parts within each stratum. First, a 0.05 percent stratified Bernoulli sample of approximately 65,000 returns was selected, called the Continuous Work History Sample (CWHS, Weber 2004). A Bernoulli sample was also selected independently from each stratum, with rates ranging from 0.01 to 100 percent. The full sample thus consists of the CWHS plus all additional returns selected via the Bernoulli sample. For Tax Year 2005, the sampling rates were increased and ranged from 0.05 to 100 percent. Table 2 contains the sample and estimated population sizes (using the sample weights, both after the sample was selected and those numbers realized after SOI’s data capture and cleaning).

Table 2: Realized Sample Sizes and Estimated Population Sizes, Tax Years 2004 and 2005

	Tax Year 2004	Tax Year 2005
<i>Sample Sizes</i>		
Originally selected	200,778	292,966
After data cleaning	200,295	292,837
<i>Estimated Population Sizes</i>		
Originally selected	133,189,982	134,494,440
After data cleaning	132,226,042	134,372,678

Every year, using condition (2.1) for every tax return automatically accounts for births, deaths, and the stratum jumpers in the population as follows:

- *Births*: each birth is independently assigned a PRN; if (2.1) holds, then the unit is selected for the sample.
- *Deaths*: units are not present in the population file, so they are not in the sample.
- *Stratum jumpers*: if, from year t_1 to t_2 , a return switches from stratum h_1 to stratum h_2 , then the return is in the sample in both years if $r_i < \min(\pi_{h_1}, \pi_{h_2})$ (i.e., if the PRN is less than the rates for both strata).

There are four conditions related to whether or not a stratum jumper is selected for the t_2 sample, which are shown in Table 3. They depend on the size of the two years’ sampling fractions (relative to each other) and if the unit was in the year t_1 sample.

Table 3: Sample Inclusion of Stratum Jumpers in Year t_2

	Was unit i in the sample in year t_1 ?	
<i>Sampling rate relationship</i>	Yes	No
$\pi_{h_1} \leq \pi_{h_2}$	Yes	Maybe
$\pi_{h_1} > \pi_{h_2}$	Maybe	No

The Table 3 conditions are further explained, where h_1 denotes the year t_1 stratum and h_2 for year t_2 . These probabilities are needed for subsequent variance calculations.

Returns selected in year t_1 :

- If $\pi_{h_1} \leq \pi_{h_2}$, then unit i will automatically be included in the year t_2 sample with conditional probability of selection 1 and unconditional probability of selection π_{h_2} , since $r_i < \pi_{h_1} \leq \pi_{h_2}$.
- If unit i switches to a stratum with a smaller sampling rate, then it will be in the year t_2 sample if $r_i < \pi_{h_2}$ with conditional probability π_{h_2}/π_{h_1} (and not in the sample with probability $1 - \pi_{h_2}/\pi_{h_1}$).

Returns not selected in year t_1 :

- If $\pi_{h_1} \leq \pi_{h_2}$, then a unit will only be in the year t_2 sample if $\pi_{h_1} < r_i \leq \pi_{h_2}$ with conditional probability $(\pi_{h_2} - \pi_{h_1})/(1 - \pi_{h_1})$ (and not in with probability $1 - (\pi_{h_2} - \pi_{h_1})/(1 - \pi_{h_1})$).
- If unit i switches to a stratum with a smaller sampling rate, then it will not be sampled at time t_2 since $\pi_{h_2} < \pi_{h_1} \leq r_i$ does not meet condition (2.1).

This sample selection method also ensures a large overlap between two years, since a sampled unit is selected in both years if $r_i \leq \pi_{h_1} \leq \pi_{h_2}$. There are a small number of 1040 tax returns where r_i changes, even though the composition of the return itself remains the same (which can occur, for example, on a married joint tax return that switches the primary and secondary SSN's, since r_i is calculated using the primary SSN). The overlapping of units across different year's samples also creates a large covariance term that must be accounted for in variance estimation of the difference between two years' estimates.

There are additional sample selection issues due to changes in population units that affect the covariance term. For example, in the 1040 sample, "marriages" occur when two tax returns that previously filed separately file as a joint-married return the next year and "divorces" when a joint married tax return becomes two separate entities (which includes both legally divorced taxpayers and married taxpayers who choose to file their returns separately). We use the following rules in the covariance estimation, which generally lead to underestimating it:

- *Marriages*: two "single" returns (filing either as single or married separate) in year t_1 that file as a joint married return are considered two deaths in year t_1 and a birth in year t_2 .
- *Divorces*: a married joint return that becomes two single entities is considered a death in year t_1 and two births in year t_2 .
- *SSN swapping*: joint married tax returns that are in both years are tracked and considered the same unit in both years.

Sample design changes also result in sampling rate changes between years. For example, the 1040 samples included returns selected for the CWSH part of the sample using certain endings of the SSN. For Tax Year 2004, only half of the possible SSN endings were edited and used to produce the 2004 cross-sectional estimates and a defacto 100 percent increase in sampling rates. However, all selected CWSH returns were edited for the 2005 sample, resulting in approximately 60,000 additional returns used to produce the 1040 cross-sectional sample estimates. Also, the Congressionally-mandated five-year Foreign Income study was selected in the 2006 sample, resulting in sampling rate increases to include approximately 15,000 more returns in the associated strata. Our estimators can account for such changes.

3. Estimators for Totals and Their Change

3.1: Notation and Probabilities of Selection

A Bernoulli sample is selected within each stratum as described in Section 3, where π_h , the stratum sampling rate in a given year, is also the probability of selection for all units in stratum h . The following random variables denote sample inclusion of unit i at times t_1 and t_2 :

$$\delta_i(t_1) = \begin{cases} 1 & \text{if unit } i \in s_1 \\ 0 & \text{otherwise} \end{cases} \quad \text{and} \quad \delta_i(t_2) = \begin{cases} 1 & \text{if unit } i \in s_2 \\ 0 & \text{otherwise} \end{cases}.$$

From these expressions, the conditional and unconditional probabilities of selection by population domain can be derived. For Bernoulli sampling, the expected values and variances of the inclusion indicator for each year are

$$E[\delta_i(t_1)] = \pi_{h_1}, \quad \text{Var}[\delta_i(t_1)] = \pi_{h_1}(1 - \pi_{h_1})$$

$$E[\delta_i(t_2)] = \pi_{h_2}, \quad \text{Var}[\delta_i(t_2)] = \pi_{h_2}(1 - \pi_{h_2}).$$

To compute $E[\delta_i(t_2)\delta_i(t_1)] - E[\delta_i(t_2)]E[\delta_i(t_1)]$ for the covariance, note that $\delta_i(t_2)\delta_i(t_1) = 1$ only when a unit is in the sample for both time periods. Thus, the covariance for the indicator variable for unit i in stratum h_1 at time t_1 and in stratum h_2 at time t_2 is given by:

$$\text{Cov}[\delta_i(t_2), \delta_i(t_1)] = \min(\pi_{h_1}, \pi_{h_2}) - \pi_{h_1}\pi_{h_2} = \Delta_{h_1h_2}.$$

3.2: Finite Population Totals of Interest

The finite population totals of a study variable of interest y at times t_1 and t_2 are denoted by

$$T(t_1) = \sum_{h_1=1}^{H_1} \sum_{i \in U_{h_1 \bullet}} y_{1i} \text{ and } T(t_2) = \sum_{h_2=1}^{H_2} \sum_{i \in U_{\bullet h_2}} y_{2i} \quad (3.1)$$

where y_{1i} and y_{2i} are the y -values (for the same variable of interest) for unit i at times t_1 and t_2 .

3.3. The Horvitz-Thompson Estimator

The Horvitz-Thompson (HT) estimators for $T(t_1), T(t_2)$ are

$$\hat{T}_{\pi}(t_1) = \sum_{h_1=1}^{H_1} \sum_{i \in U_{h_1 \bullet}} \frac{\delta_i(t_1)y_{1i}}{\pi_{h_1}} = \sum_{h_1=1}^{H_1} \hat{T}_{h_1 \bullet} \text{ and } \hat{T}_{\pi}(t_2) = \sum_{h_2=1}^{H_2} \sum_{i \in U_{\bullet h_2}} \frac{\delta_i(t_2)y_{2i}}{\pi_{h_2}} = \sum_{h_2=1}^{H_2} \hat{T}_{\bullet h_2}, \quad (3.2)$$

where $1/\pi_{h_1}$ and $1/\pi_{h_2}$ are the base weights for unit i at times t_1 and t_2 and $\hat{T}_{h_1 \bullet} = 1/\pi_{h_1} \sum_{i \in U_{h_1 \bullet}} \delta_i(t_1)y_{1i}$, $\hat{T}_{\bullet h_2} = 1/\pi_{h_2} \sum_{i \in U_{\bullet h_2}} \delta_i(t_2)y_{2i}$ are HT estimators for each year's stratum totals. The estimators $\hat{T}_{\pi}(t_1)$ and $\hat{T}_{\pi}(t_2)$ are unconditionally (i.e., over all possible samples for each year) unbiased for $T(t_1)$ and $T(t_2)$, respectively.

3.4. The Poststratified (Conditional HT) Estimator

Even though the HT estimator is unconditionally unbiased for the population total, it can have a high variance since the sample size is a random variable under Bernoulli sampling. Instead, SOI uses a poststratified (PS) estimator that conditions on the number of achieved units in each stratum. This estimator, which is conditionally unbiased for the population total (Brewer *et al.*, 1972), reduces the variability caused by the random stratum sample sizes and leads to formulae simplifications.

First, the observed number of sample returns in stratum h from year t_1 is denoted by $n_{h_1 \bullet} = \sum_{i \in U_{h_1 \bullet}} \delta_i(t_1)$. Assuming that $n_{h_1 \bullet} > 0$, it can be shown that conditional on $\{n_{1 \bullet}, n_{2 \bullet}, \dots, n_{H_1 \bullet}\}$, the sample design at time t_1 is a stratified simple random sample with stratum sample sizes $n_{1 \bullet}, n_{2 \bullet}, \dots, n_{H_2 \bullet}$. Thus, for $N_{h_1 \bullet}$ denoting the number of population units in stratum h_1 at time t_1 , the (conditional) HT estimator for $T(t_1)$ is

$$\hat{T}_{\pi c}(t_1) = \sum_{h_1=1}^{H_1} \frac{N_{h_1 \bullet}}{n_{h_1 \bullet}} \sum_{i \in U_{h_1 \bullet}} \delta_i(t_1)y_{1i}. \quad (3.3)$$

Similarly, if $n_{\bullet h_2} = \sum_{i \in U_{\bullet h_2}} \delta_i(t_2) > 0$, then conditional on $\{n_{\bullet 1}, n_{\bullet 2}, \dots, n_{\bullet H_2}\}$, the sample design at time t_2 is a stratified simple random sample with stratum sample sizes $n_{\bullet 1}, n_{\bullet 2}, \dots, n_{\bullet H_2}$. For $N_{\bullet h_2}$ being the number of population elements in stratum h at time t_2 , the (conditional) HT estimator for $T(t_2)$ is

$$\hat{T}_{\pi c}(t_2) = \sum_{h_2=1}^{H_2} \frac{N_{\bullet h_2}}{n_{\bullet h_2}} \sum_{i \in U_{\bullet h_2}} \delta_i(t_2)y_{2i}. \quad (3.4)$$

SOI uses $\hat{T}_{\pi c}(t_1)$ and $\hat{T}_{\pi c}(t_2)$ to estimate time-specific totals, which are conditionally unbiased for their corresponding population totals. They are special forms of the PS estimator, where the poststrata are the same as the design strata. The PS estimators (which are not unconditionally unbiased for the population totals, since they involve a ratio) for times t_1 and t_2 are

$$\hat{T}_{PS}(t_1) = \sum_{h_1=1}^{H_1} \frac{N_{h_1 \bullet}}{\hat{N}_{h_1 \bullet}} \hat{T}_{h_1 \bullet} \text{ and } \hat{T}_{PS}(t_2) = \sum_{h_2=1}^{H_2} \frac{N_{\bullet h_2}}{\hat{N}_{\bullet h_2}} \hat{T}_{\bullet h_2}, \quad (3.5)$$

where $\hat{N}_{h_1 \bullet} = \frac{1}{\pi_{h_1}} \sum_{i \in U_{h_1 \bullet}} \delta_i(t_1)$ is the estimated number of stratum h_1 population units for year 1. For year 2, $\hat{N}_{\bullet h_2}$ is similarly defined. It can be shown, by definition of $\hat{T}_{h_1 \bullet}$ and $\hat{T}_{\bullet h_2}$, that $\hat{T}_{PS}(t_1)$ reduces to $\hat{T}_{\pi c}(t_1)$ and $\hat{T}_{PS}(t_2)$ reduces to $\hat{T}_{\pi c}(t_2)$.

3.5. Estimators of Change

The change in level between two time points is denoted by

$$D = T(t_2) - T(t_1). \quad (3.6)$$

The conditional and unconditional estimators of time-specific totals lead to two estimators of this difference. Based on the

unconditional HT estimators, we have

$$\hat{D}_\pi = \hat{T}_\pi(t_2) - \hat{T}_\pi(t_1), \quad (3.7)$$

which is unconditionally unbiased for the change in level between time t_1 and t_2 . By breaking $\hat{T}_\pi(t_1)$ into the sum of births for time t_2 and units in both year's sample summed over the year 1 strata, and $\hat{T}_\pi(t_2)$ into the sum of the deaths for time t_1 and the units in both samples over the year 2 strata, expression (3.7) can be rewritten as:

$$\hat{D}_\pi = \sum_{h_2=1}^{H_2} \sum_{i \in U_{0h_2}} \frac{\delta_i(t_2)y_{2i}}{\pi_{h_2}} - \sum_{h_1=1}^{H_1} \sum_{i \in U_{h_10}} \frac{\delta_i(t_1)y_{1i}}{\pi_{h_1}} + \sum_{h_1=1}^{H_1} \sum_{h_2=1}^{H_2} \sum_{i \in U_{h_1h_2}} \left[\frac{\delta_i(t_2)y_{2i}}{\pi_{h_2}} - \frac{\delta_i(t_1)y_{1i}}{\pi_{h_1}} \right] \quad (3.8)$$

Similarly, based on the conditional HT estimators,

$$\hat{D}_{PS} = \hat{T}_{PS}(t_2) - \hat{T}_{PS}(t_1), \quad (3.9)$$

which is conditionally unbiased for the change in level. Similarly, this estimator can be rewritten as:

$$\hat{D}_{PS} = \sum_{h_2=1}^{H_2} \sum_{i \in U_{0h_2}} \frac{N_{\bullet h_2} \delta_i(t_2)y_{2i}}{n_{\bullet h_2}} - \sum_{h_1=1}^{H_1} \sum_{i \in U_{h_10}} \frac{N_{h_1 \bullet} \delta_i(t_1)y_{1i}}{n_{h_1 \bullet}} + \sum_{h_1=1}^{H_1} \sum_{h_2=1}^{H_2} \sum_{i \in U_{h_1h_2}} \left[\frac{N_{\bullet h_2} \delta_i(t_2)y_{2i}}{n_{\bullet h_2}} - \frac{N_{h_1 \bullet} \delta_i(t_1)y_{1i}}{n_{h_1 \bullet}} \right] \quad (3.10)$$

4. Theoretical Variances

4.1: Single Year Variances

4.1.1: Unconditional HT Estimators

Using standard Poisson sampling results (Result 3.2.1 from Sarndal *et al.*, (1992)), the variances of the unconditional HT estimators for times t_1 and t_2 are

$$Var[\hat{T}_\pi(t_1)] = \sum_{h_1=1}^{H_1} \frac{(1-\pi_{h_1})}{\pi_{h_1}} \sum_{i \in U_{h_1 \bullet}} y_{1i}^2 \quad (4.1)$$

$$Var[\hat{T}_\pi(t_2)] = \sum_{h_2=1}^{H_2} \frac{(1-\pi_{h_2})}{\pi_{h_2}} \sum_{i \in U_{\bullet h_2}} y_{2i}^2 \quad (4.2)$$

4.1.2: Unconditional Variance of PS Estimators

Using $N_h(1/\pi_h - 1) \approx N_h^2/n_h(1 - n_h/N_h)$, Sarndal *et al.* (expression 3.2.7, p. 65) approximated the unconditional variance of the PS estimators. Their approximation, adjusted for our notation, is

$$Var[\hat{T}_{PS}(t_1)] \approx \sum_{h_1=1}^{H_1} N_{h_1 \bullet} (1/\pi_{h_1} - 1) S_{h_1 \bullet}^2 \quad (4.3)$$

$$Var[\hat{T}_{PS}(t_2)] \approx \sum_{h_2=1}^{H_2} N_{\bullet h_2} (1/\pi_{h_2} - 1) S_{\bullet h_2}^2, \quad (4.4)$$

where $S_{h_1 \bullet}^2 = 1/(N_{h_1 \bullet} - 1) \sum_{i \in U_{h_1 \bullet}} (y_{1i} - \bar{Y}_{h_1 \bullet})^2$ and $S_{\bullet h_2}^2 = 1/(N_{\bullet h_2} - 1) \sum_{i \in U_{\bullet h_2}} (y_{2i} - \bar{Y}_{\bullet h_2})^2$ are the population stratum variances for t_1 and t_2 and the population strata means are $\bar{Y}_{h_1 \bullet} = 1/N_{h_1 \bullet} \sum_{i \in U_{h_1 \bullet}} y_{1i}$ and $\bar{Y}_{\bullet h_2} = 1/N_{\bullet h_2} \sum_{i \in U_{\bullet h_2}} y_{2i}$.

4.1.3: Conditional Variances of PS Estimators

Holt and Smith (1979) observed that conditioning on an achieved post stratum sample size is inferentially more appropriate than averaging over all possible sample sizes, as in (4.1) and (4.2). The theoretical conditional variances of the PS estimators for both years are simply the variances of a total under stratified simple random sampling:

$$Var[\hat{T}_{PS}(t_1)] = \sum_{h_1=1}^{H_1} N_{h_1 \bullet}^2 / n_{h_1 \bullet} (1 - n_{h_1 \bullet} / N_{h_1 \bullet}) S_{h_1 \bullet}^2 \quad (4.5)$$

$$Var[\hat{T}_{PS}(t_2)] = \sum_{h_2=1}^{H_2} N_{\bullet h_2}^2 / n_{\bullet h_2} (1 - n_{\bullet h_2} / N_{\bullet h_2}) S_{\bullet h_2}^2 \quad (4.6)$$

The variances in (4.5) and (4.6) are preferable since they reflect the precision for the sample sizes actually obtained. In comparing expressions (4.1) to (4.5) and (4.2) to (4.6), in almost all practical situations, the conditional variances of the PS estimators are substantially smaller than the unconditional variances of the HT estimators. To see this, in general, we can write (expanding Exp.3.2.5 in Sarndal *et al.* (1992) to stratified sampling):

$$\begin{aligned} \text{Var}_{STBE} [\hat{T}_\pi] &= \text{Var}_{STSI} [\hat{T}_\pi] + \sum_{h=1}^H N_h^2/n_h (1-n_h/N_h) S_h^2 \left[1-1/N_h + 1/CV_{yh}^2 \right] \\ &\approx \text{Var}_{STSI} [\hat{T}_\pi] + \sum_{h=1}^H N_h (N_h/n_h - 1) \bar{Y}_h^2 \left[1 + CV_{yh}^2/N_h \right] \end{aligned} \quad (4.7)$$

Thus, the theoretical variance of a total under stratified Bernoulli sampling is equal to the stratified simple random sampling variance plus an additional factor (of the same magnitude) that depends on the stratum population mean of the study variable.

4.2: Variance of the Difference

4.2.1: Unconditional Variance of the HT Estimators

For the unconditional HT estimators, it can be shown that the unconditional variance of the difference is the variance from the year 1, births plus the variance from the year 2 deaths, plus the variance from returns in the same or different strata in both years. In formula form, this is:

$$\text{Var} [\hat{D}_\pi] = \sum_{h_2=1}^{H_2} \frac{1-\pi_{h_2}}{\pi_{h_2}} \sum_{i \in U_{\bullet h_2}} y_{2i}^2 + \sum_{h_1=1}^{H_1} \frac{1-\pi_{h_1}}{\pi_{h_1}} \sum_{i \in U_{h_1 \bullet}} y_{1i}^2 + \sum_{h_1=1}^{H_1} \sum_{h_2=1}^{H_2} \sum_{i \in U_{h_1 h_2}} \left[\frac{(1-\pi_{h_2}) y_{2i}^2}{\pi_{h_2}} + \frac{(1-\pi_{h_1}) y_{1i}^2}{\pi_{h_1}} - 2 \frac{\Delta_{h_1 h_2} y_{2i} y_{1i}}{\pi_{h_1} \pi_{h_2}} \right] \quad (4.8)$$

4.2.2: Unconditional Variance of the PS Estimators (Linear Approximation)

Using linear approximations to the PS estimators, the unconditional variance of the difference is

$$\text{Var} [\hat{D}_{PS}] \approx \sum_{h_2=1}^{H_2} \frac{1-\pi_{h_2}}{\pi_{h_2}} N_{\bullet h_2} S_{\bullet h_2}^2 + \sum_{h_1=1}^{H_1} \frac{1-\pi_{h_1}}{\pi_{h_1}} N_{h_1 \bullet} S_{h_1 \bullet}^2 - 2 \sum_{h_1=1}^{H_1} \sum_{h_2=1}^{H_2} \left[\frac{\Delta_{h_1 h_2} N_{h_1 h_2} S_{h_1 h_2}}{\pi_{h_1} \pi_{h_2}} \right] \quad (4.9)$$

where $S_{h_1 h_2} = 1/(N_{h_1 h_2} - 1) \sum_{i \in U_{h_1 h_2}} (y_{2i} - \bar{Y}_{\bullet h_2})(y_{1i} - \bar{Y}_{h_1 \bullet})$.

4.2.3: Unconditional Variance of PS Estimators (Substitution Form)

The variance in (4.8) can be expressed in a more standard form by converting some of the summations into stratum variances and covariances and approximating the sampling rates using the actual sample and population sizes achieved in each stratum. One approach is substituting marginal actual sampling rates for terms like $1-\pi_h$ and replace $1/\pi_h$ by a stratum population size divided by the actual stratum sample size. Using these, we get

$$\text{Var} [\hat{D}_{PS}] \approx \sum_{h_2=1}^{H_2} \left(1 - \frac{n_{\bullet h_2}}{N_{\bullet h_2}} \right) \frac{N_{\bullet h_2}^2}{n_{\bullet h_2}} S_{\bullet h_2}^2 + \sum_{h_1=1}^{H_1} \left(1 - \frac{n_{h_1 \bullet}}{N_{h_1 \bullet}} \right) \frac{N_{h_1 \bullet}^2}{n_{h_1 \bullet}} S_{h_1 \bullet}^2 - 2 \sum_{h_1=1}^{H_1} \sum_{h_2=1}^{H_2} \left[\frac{\Delta_{h_1 h_2} N_{h_1 h_2} S_{h_1 h_2}}{\pi_{h_1} \pi_{h_2}} \right] \quad (4.10)$$

5. Variance Estimators

5.1: Single-Year Variance Estimators

We assume that the terms $N_{h_1 \bullet}$ and $N_{\bullet h_2}$ are known for all h_1, h_2 and consider both cases of the $N_{h_1 h_2}$ being known and unknown. Assuming that the $N_{h_1 h_2}$ are unknown, these are conditionally unbiased estimators of the strata population means:

$$\bar{y}_{h_1 \bullet} = 1/n_{h_1 \bullet} \sum_{i \in U_{h_1 \bullet}} \delta_i(t_1) y_{1i} \quad \text{and} \quad \bar{y}_{\bullet h_2} = 1/n_{\bullet h_2} \sum_{i \in U_{\bullet h_2}} \delta_i(t_2) y_{2i} .$$

Since conditionally (and approximately unconditionally) unbiased estimators for the strata variances $S_{h_1 \bullet}^2$ and $S_{\bullet h_2}^2$ are $s_{h_1 \bullet}^2 = 1/(n_{h_1 \bullet} - 1) \sum_{i \in U_{h_1 \bullet}} \delta_i(t_1) (y_{1i} - \bar{y}_{h_1 \bullet})^2$ and $s_{\bullet h_2}^2 = 1/(n_{\bullet h_2} - 1) \sum_{i \in U_{\bullet h_2}} \delta_i(t_2) (y_{2i} - \bar{y}_{\bullet h_2})^2$, the within-year variance estimators are the standard variance estimators under stratified simple random sampling:

$$\text{var} [\hat{T}_{PS}(t_1)] = \sum_{h_1=1}^{H_1} N_{h_1 \bullet}^2 / n_{h_1 \bullet} (1 - n_{h_1 \bullet} / N_{h_1 \bullet}) s_{h_1 \bullet}^2 \quad (5.1)$$

$$\text{var} [\hat{T}_{PS}(t_2)] = \sum_{h_2=1}^{H_2} N_{\bullet h_2}^2 / n_{\bullet h_2} (1 - n_{\bullet h_2} / N_{\bullet h_2}) s_{\bullet h_2}^2 \quad (5.2)$$

These are conditionally unbiased for (4.5) and (4.6).

5.2: Variance Estimators of the Differences

5.2.1: The Unconditional Difference

Using sample-based estimates for each (4.8) component, we get the following estimate of $\text{Var} [\hat{D}_\pi]$:

$$var[\hat{D}_\pi] = \sum_{h_2=1}^{H_2} \frac{1-\pi_{h_2}}{\pi_{h_2}^2} \sum_{i \in s_{\bullet h_2}} y_{2i}^2 + \sum_{h_1=1}^{H_1} \frac{1-\pi_{h_1}}{\pi_{h_1}^2} \sum_{i \in s_{h_1 \bullet}} y_{1i}^2 - 2 \sum_{h_1=1}^{H_1} \sum_{h_2=1}^{H_2} \sum_{i \in s_{h_1 h_2}} \frac{\Delta_{h_1 h_2} y_{2i} y_{1i}}{\min(\pi_{h_1}, \pi_{h_2}) \pi_{h_1} \pi_{h_2}} \quad (5.3)$$

5.2.2: The Conditional Difference

Using sample-based estimates for each (4.10) component, we have the approximate estimator of $Var[\hat{D}_{PS}]$:

$$var[\hat{D}_{PS}] \approx \sum_{h_2=1}^{H_2} \left(1 - \frac{n_{\bullet h_2}}{N_{\bullet h_2}}\right) \frac{N_{\bullet h_2}^2}{n_{\bullet h_2}} s_{\bullet h_2}^2 + \sum_{h_1=1}^{H_1} \left(1 - \frac{n_{h_1 \bullet}}{N_{h_1 \bullet}}\right) \frac{N_{h_1 \bullet}^2}{n_{h_1 \bullet}} s_{h_1 \bullet}^2 - 2 \sum_{h_1=1}^{H_1} \sum_{h_2=1}^{H_2} \frac{\Delta_{h_1 h_2} \hat{N}_{h_1 h_2} c_{h_1 h_2}}{\pi_{h_1} \pi_{h_2}} \quad (5.4)$$

where $\hat{N}_{h_1 h_2} = n_{h_1 h_2} / \min(\pi_{h_1}, \pi_{h_2})$, $\pi_{h_2} = n_{\bullet h_2} / N_{\bullet h_2}$, $\pi_{h_1} = n_{h_1 \bullet} / N_{h_1 \bullet}$, and the *unweighted* covariance between the y -values for units in both years' samples is $c_{h_1 h_2} = 1 / (n_{h_1 h_2} - 1) \sum_{i \in s_{h_1 h_2}} (y_{2i} - \bar{y}_{\bullet h_2})(y_{1i} - \bar{y}_{h_1 \bullet})$. Note that the (5.4) covariance term is an estimator of the unconditional covariance, but it does partially account for achieved sample sizes in using $n_{h_1 \bullet} / N_{h_1 \bullet}$ and $n_{\bullet h_2} / N_{\bullet h_2}$ in place of π_{h_1} and π_{h_2} .

6. Data Elements Required For Variance Estimators

The following information is needed to evaluate (5.3) and (5.4):

- The strata occupied by each sample unit at times t_1 and t_2 and the data values for every unit in the sample each time, y_{1i} and y_{2i} .
- $N_{h_1 \bullet}$, the population size of stratum h_1 at time t_1 and $N_{\bullet h_2}$ the population size of stratum h_2 at time t_2 .
- π_{h_1} , the probability of selection for a unit in stratum h_1 at time t_1 , and π_{h_2} , the selection probability for a unit in stratum h_2 at time t_2 .
- $n_{h_1 h_2}$, $N_{h_1 h_2}$ and $c_{h_1 h_2}$, the number of sample and population units and unweighted covariance in each of the $(h_1 h_2)$ cells.

To account for SOI's sample including prior year returns in each sample, we matched the most recent tax return within each year together in both the population and sample. This led to ignoring a few cases where a taxpayer filed more than one return in a single year even though these returns were used in estimating totals. Doing so led to a slight underestimation of the covariances, but the impact of this was much less than ignoring the covariance term.

7. Results

Table 5 shows the ratios of the unconditional to conditional estimates for each year's estimated total, the difference between them, and the associated variance estimates for eight variables estimated from SOI's Tax Year 2004 and 2005 Individual samples. While the point estimates in Table 5 are essentially identical, the unconditional variances are much larger (as much as six to ten times larger, in Adjusted Gross Income and Taxable Income) than the corresponding conditional variance estimates when the means are larger for each year, as expected from (4.7).

Table 5. Single-Year and Between-Year Difference Estimates, by Tax Year and Variable of Interest

Variable	Single-Year Estimates						Between-Year Difference Estimates		
	Estimated Means (in \$'s)		HT Total / PS Total		HT Variance / PS Variance		HT Total / PS Total	HT Variance / PS Variance*	
	2004	2005	2004	2005	2004	2005	(2005-2004)	(5.4) v1	(5.4) v2
Adjusted Gross Income	51,342	55,238	0.994	1.001	10.638	9.739	1.074	9.609	9.737
Taxable Income	35,320	38,231	0.994	1.001	6.642	6.157	1.066	6.347	6.448
Total Income Tax	6,292	6,957	0.995	1.000	3.891	3.827	1.041	4.322	4.419
Business or profession net income (less loss)	1,870	2,007	1.004	1.007	1.278	1.306	1.042	1.801	1.887
Alternative minimum tax	99	130	1.002	1.001	1.194	1.218	0.999	1.416	1.426
Net capital gain (less loss)	3,568	4,934	1.001	1.001	1.175	1.225	1.001	1.535	1.568
Charitable contributions	1,252	1,365	0.996	1.001	1.272	1.270	1.051	1.934	2.066
Charitable contributions other than cash	328	358	0.997	1.001	1.015	1.008	1.037	1.087	1.098

* v1 is expression (5.4) using the estimated $\hat{N}_{h_1 h_2}$ counts; v2 is (5.4) using the known $N_{h_1 h_2}$ counts.

Table 6 contains the ratio of the variance estimates of the unconditional and conditional differences in the 2004 and 2005 totals for our eight variables, when estimating and ignoring the covariance term in (5.3) and (5.4). We also include results when estimating the $N_{h_1 h_2}$ or using the known counts in (5.4). Ignoring the covariance leads to excessively large variance estimates because the benefit of having a large sample overlap is ignored. For example, ignoring the covariance would result

in an estimated variance of the difference in HT estimators that was 68% too large and 57% to 59% too large for the variance of the difference in the PS estimators. While estimating the $N_{h_1h_2}$ did not lead to much different covariance estimates for these national-level variables, the HT estimates had higher decreases in the variance of the difference due to the estimated covariances for all variables except net capital gain (less loss), where the percentages were close. We also calculated the t -statistics to test whether the population differences were zero. While the t -statistics had different values depending on which variance estimator was used, all were highly significant, and were thus omitted.

Table 6. Ratios of Variances of Differences when Ignoring the Covariance to Variances When Estimating It

Variable	HT Difference	PS Difference (Using Estimated $\hat{N}_{h_1h_2}$)	PS Difference (Using Known $N_{h_1h_2}$)
Adjusted Gross Income	1.682	1.574	1.595
Taxable Income	1.660	1.636	1.662
Total Income Tax	1.648	1.844	1.885
Business or profession net income (less loss)	1.537	2.145	2.248
Alternative minimum tax	1.242	1.456	1.466
Net capital gain (less loss)	1.099	1.403	1.433
Charitable contributions	1.546	2.352	2.512
Charitable contributions other than cash	1.077	1.158	1.169

8. Conclusions

The large overlap of units between SOI's 2004 and 2005 Individual tax return samples resulted in a large covariance term in both the conditional and unconditional variances. In comparing the (5.3) and (5.4) estimates, using the unconditional (5.3) formulas produced larger estimates for the separate year variances. This is due to the unconditional formulas incorporating the extra variability due to random sample sizes. The most interesting result was that these variances are much larger than the associated PS estimates of the single year totals, despite the fact the HT and PS point estimates of totals are almost identical. The SOI data support expression (4.7), that is, the mean of the variable of interest affects the size of the variance estimate. Since valid inferences are obtained using the conditional variance estimates, in summary, the best estimation strategy is using expression (5.4) to estimate the variance of the difference in PS estimators with the known $N_{h_1\bullet}$, $N_{\bullet h_2}$, and $N_{h_1h_2}$ counts.

Despite large computing resources needed to match the two year's population files, it was not difficult to compute the estimates once the $n_{h_1h_2}$, $N_{h_1h_2}$ and $c_{h_1h_2}$ quantities were produced. Notably, after the two population files were merged, only the $N_{h_1h_2}$ counts were needed. A more complicated estimator of between-year change, such as a relative change in the two years' estimated totals, would require a more sophisticated approach (such as the Taylor series approximation used in Berger 2004 and Nordberg 2000). Our estimators would also need to be slightly modified for domain estimation. Such extensions are future consideration topics.

9. References

- Berger, Y.G. (2004), "Variance Estimation for Measures of Change in Probability Sampling," *The Canadian Journal of Statistics*, 32, 451-467.
- Brewer, K.R.W., Early, L.J., and Joyce, S.F. (1972), "Selecting Several Samples from a Single Population," *Australian Journal of Statistics*, 14, 231-239.
- Cooney, S. (2008), Personal communication.
- Holt, D. and Smith, T.M.F. (1979), "Post Stratification," *Journal of the Royal Statistical Society A*, 142, 33-46.
- Nordberg, L. (2000) "On Variance Estimation for Measures of Change When Samples are Coordinated by the Use of Permanent Random Numbers," *Journal of Official Statistics*, 14, No. 4, 363-368.
- Sarndal, Swensson, and Wretman (1992), *Model Assisted Survey Sampling*, Springer-Verlag, New York.
- Testa, V. and Scali, J. (2006), *Statistics of Income—2004 Individual Income Tax Returns, IRS, Publication 1304*, 23-27.
- Valliant, R. and Casady, R., (1998), "The Variance of An Estimator of Change From the Sample of Partnership Tax Returns," *Statistical Issues Related to SOI Family Cross-Sectional and Longitudinal Studies*, Contract No. TIRN0-96-D-0030 Task 5, Washington DC.
- Weber, M., (2004), "The Statistics of Income 1979-2002 Continuous Work History Sample Individual Tax Return Panel," <http://www.irs.gov/pub/irs-soi/04webasa.pdf>.
- Wood, J. (2008). "On the Covariance Between Related Horvitz-Thompson Estimators," *Journal of Official Statistics*, 24, No. 1, 53-78.